

The Search for Higher Helicities

Clayton Shonkwiler

Department of Mathematics
Haverford College

January 8, 2011

Joint work with: Dennis DeTurck, Herman Gluck, Rafal Komendarczyk, Paul Melvin, and David Shea Vela-Vick

Let V be a vector field on a compact domain $\Omega \subset \mathbb{R}^3$.

The *energy* of V is

$$E(V) = \int_{\Omega} V \cdot V \, \text{dvol}_{\Omega}.$$

Let V be a vector field on a compact domain $\Omega \subset \mathbb{R}^3$.

The *energy* of V is

$$E(V) = \int_{\Omega} V \cdot V \, \text{dvol}_{\Omega}.$$

Suppose V is divergence-free. Then

$$V = \nabla \times W,$$


where W is the vector potential for V .

If V is a plasma flow, is there any obstruction to V relaxing to have arbitrarily small energy?


If V is a plasma flow, is there any obstruction to V relaxing to have arbitrarily small energy?

Mathematically, are there obstructions to making the field energy arbitrarily small via volume-preserving diffeomorphisms homotopic to the identity?


Answer: Not necessarily!




A field which can be relaxed to have arbitrarily small energy (Freedman).



Linked orbits prevent the energy from getting arbitrarily small.



$$1 = Lk(K, L) = \frac{1}{4\pi} \int_{K \times L} \frac{dx}{ds} \times \frac{dy}{dt} \cdot \frac{x - y}{|x - y|^3} ds dt$$



$$2 = Lk(K, L) = \frac{1}{4\pi} \int_{K \times L} \frac{dx}{ds} \times \frac{dy}{dt} \cdot \frac{x - y}{|x - y|^3} ds dt$$

The Gauss Linking Integral:

$$Lk(K, L) = \frac{1}{4\pi} \int_{K \times L} \frac{dx}{ds} \times \frac{dy}{dt} \cdot \frac{x - y}{|x - y|^3} ds dt$$

The Gauss Linking Integral:

$$Lk(K, L) = \frac{1}{4\pi} \int_{K \times L} \frac{dx}{ds} \times \frac{dy}{dt} \cdot \frac{x - y}{|x - y|^3} ds dt$$

Important: the integrand is isometry-invariant.

The Gauss Linking Integral:

$$Lk(K, L) = \frac{1}{4\pi} \int_{K \times L} \frac{dx}{ds} \times \frac{dy}{dt} \cdot \frac{x - y}{|x - y|^3} ds dt$$

Important: the integrand is isometry-invariant.

Definition (Woltjer)

The *helicity* of a divergence-free vector field V on a compact domain $\Omega \subset \mathbb{R}^3$ is

$$H(V) := \frac{1}{4\pi} \int_{\Omega \times \Omega} V(x) \times V(y) \cdot \frac{x - y}{|x - y|^3} d\text{vol}_x d\text{vol}_y.$$

Theorem

$H(V)$ is invariant under any volume-preserving diffeomorphism which is homotopic to the identity.

Theorem (Arnol'd)


$$|H(V)| \leq R(\Omega)E(V)$$

where $R(\Omega)$ is a positive constant depending on the shape and size of Ω .

Arnol'd and Khesin:


The dream is to define such a hierarchy of invariants for generic vector fields such that, whereas all the invariants of order $\leq k$ have zero value for a given field and there exists a nonzero invariant of order $k + 1$, this nonzero invariant provides a lower bound for the field energy.

Idea: Higher-order linking invariants



Three-component links were classified up to *link-homotopy* by Milnor:

- The pairwise linking numbers p, q, r .
- The *triple linking number* $\mu = \bar{\mu}_{123}$, which is an integer modulo $\gcd(p, q, r)$.



If V is supported on flux tubes, then μ provides a lower bound for the field energy (Monastyrsky–Retakh, Berger, etc.; see also Freedman–He).

Linking integrals and configuration spaces

Where does the Gauss Linking Integral come from?

Linking integrals and configuration spaces

Where does the Gauss Linking Integral come from?

$$S^1 \times S^1 \hookrightarrow \text{Conf}_2 \mathbb{R}^3 \xrightarrow{\simeq} S^2$$

$$(s, t) \longmapsto (x(s), y(t)) \longmapsto \frac{x(s) - y(t)}{|x(s) - y(t)|}$$

Linking integrals and configuration spaces

Where does the Gauss Linking Integral come from?

$$S^1 \times S^1 \hookrightarrow \text{Conf}_2 \mathbb{R}^3 \xrightarrow{\simeq} S^2$$

$$(s, t) \longmapsto (x(s), y(t)) \longmapsto \frac{x(s) - y(t)}{|x(s) - y(t)|}$$

If f is the composition of the above maps

$$Lk(K, L) = \int_{S^1 \times S^1} f^* \omega,$$

where ω is the standard area form on S^2 scaled to have area 1 instead of 4π .

Consider a three-component link in S^3 :

$$S^1 \times S^1 \times S^1 \hookrightarrow \text{Conf}_3 S^3 \xrightarrow{\cong} S^3 \times S^2 \xrightarrow{\pi} S^2.$$

Consider a three-component link in S^3 :

$$S^1 \times S^1 \times S^1 \hookrightarrow \text{Conf}_3 S^3 \xrightarrow{\simeq} S^3 \times S^2 \xrightarrow{\pi} S^2.$$

This determines a map

$$\begin{array}{ccc} g : & \mathcal{L}(3) & \longrightarrow [S^1 \times S^1 \times S^1, S^2] \\ & L & \longmapsto [gL] \end{array}$$

from link-homotopy classes of 3-component links to homotopy classes of maps from the 3-torus to the 2-sphere.

Consider a three-component link in S^3 :

$$S^1 \times S^1 \times S^1 \hookrightarrow \text{Conf}_3 S^3 \xrightarrow{\cong} S^3 \times S^2 \xrightarrow{\pi} S^2.$$

This determines a map

$$\begin{array}{ccc} g : & \mathcal{L}(3) & \longrightarrow [S^1 \times S^1 \times S^1, S^2] \\ & L & \longmapsto [gL] \end{array}$$

from link-homotopy classes of 3-component links to homotopy classes of maps from the 3-torus to the 2-sphere.

$[S^1 \times S^1 \times S^1, S^2]$ was classified by Pontryagin by:

- The degrees p, q, r on the 2-dimensional subtori.
- A “Hopf invariant” ν , which is an integer modulo $2 \gcd(p, q, r)$.

Interpreting link-homotopy invariants as homotopy invariants

Theorem (DGKMSV)

The map $g : \mathcal{L}(3) \rightarrow [S^1 \times S^1 \times S^1, S^2]$ is injective and maps

$$p \mapsto p$$

$$q \mapsto q$$

$$r \mapsto r$$

$$\mu \mapsto 2\mu$$

Interpreting link-homotopy invariants as homotopy invariants

Theorem (DGKMSV)

The map $g : \mathcal{L}(3) \rightarrow [S^1 \times S^1 \times S^1, S^2]$ is injective and maps

$$p \mapsto p$$

$$q \mapsto q$$

$$r \mapsto r$$

$$\mu \mapsto 2\mu$$

This implies that $\mathcal{L}(3) \rightarrow [S^1 \times S^1 \times S^1, \text{Conf}_3\mathbb{R}^3]$ is injective, proving the $n = 3$ case of a conjecture of Koschorke.

A more symmetric version of g along with a modification of Whitehead's integral for the Hopf invariant of a map $S^3 \rightarrow S^2$ yields:

Theorem (DGKMSV)

$$\mu(L) = \frac{1}{2} \int_{T^3} \delta(\varphi * \omega_L) \wedge \omega_L,$$

where $\omega_L = g_L^* \omega$ and φ is the fundamental solution of the Laplacian on T^3 .

A more symmetric version of g along with a modification of Whitehead's integral for the Hopf invariant of a map $S^3 \rightarrow S^2$ yields:


Theorem (DGKMSV)

$$\mu(L) = \frac{1}{2} \int_{T^3} \delta(\varphi * \omega_L) \wedge \omega_L,$$

where $\omega_L = g_L^* \omega$ and φ is the fundamental solution of the Laplacian on T^3 .

Note: The integrand is isometry-invariant.

The fundamental solution of the Laplacian



$$\varphi(\mathbf{x}) = \frac{1}{8\pi^3} \sum_{\mathbf{n} \in \mathbb{Z}^3 \setminus \{0\}} \frac{e^{i\mathbf{n} \cdot \mathbf{x}}}{|\mathbf{n}|^2}.$$

Thanks!

